

# The Global Land Surface Model ORCHIDEE

(ORganizing Carbon and Hydrology In Dynamic Ecosystems Environment)



**ORCHIDEE**  
LAND SURFACE MODEL

Presented by Nicolas Vuichard  
LSCE/IPSL

*Introduction - Training on ORCHIDEE model - January 2020*



ORCHIDEE  
LAND SURFACE MODEL



# Outline


---

- A brief history of ORCHIDEE & motivations
- Formalism
- Main processes
- Configurations & Inputs requirements




# A brief history

Model



(Laval et al., 1981)      (Ducoudré et al., 1993)      (Viovy et al., 1997)      (Polcher et al., 1998)      (Krinner et al., 2005)

Project / Users



# Objective

---

- Simulate Energy, Water, Carbon and Nitrogen fluxes at the land surface/atmosphere interface.
  - To be used for being the 'land surface' component of a Earth system model (IPSL-CM6).
    - Global => to represent the main vegetation cover.
    - Regional => to study feedback processes.
  - For past, present and future climates
    - Module of vegetation dynamic
    - Process-based modeling
- Conservation of mass and energy is a guiding principle for ORCHIDEE.



# Outline


---

- A brief history of ORCHIDEE & motivations
- **Formalism**
- Main processes
- Configurations & Inputs requirements



# A mosaic of vegetation and soil moisture


- Tiling for vegetation-related processes



# A mosaic of vegetation and soil moisture


## Tiling for soil hydrology

5' USDA texture map (Reynolds et al., 2000)




Dominant texture in each ORCHIDEE grid-cell:  
defining the hydraulic properties



Sub-grid scale heterogeneity:  
3 soil columns based on PFTs  
with independent water budget  
but same texture



- 1: Bare soil PFT
- 2: All Forest PFTs
- 3: All grassland and cropland PFTs



# A single energy budget



- One surface temperature per grid cell
- No vertical discretization within the canopy



# Surface variability representation ?

- In each grid cell, we account for:
  - Bare soil :  $veget\_max(1)$
  - Vegetated lands :  $veget\_max(2:nvm)$
  - Other lands (so far, only the continental ice)  
 $frac\_nobio$


$$\sum_{i=1}^{nvm} (veget\_max_i) + frac\_nobio = 1$$

- Use also of  $veget\_cov\_max = veget\_max / (1 - frac\_nobio)$
- One soil type per grid cell but different soil moisture profiles.



# Vegetated lands

- Concept of 'Plant Functional Types' (PFT)
- Defined according to systematic, physiological, phenological, climatic conditions




# Plant Functional Types


- A same set of equations governs PFT
- But parameter values differ among PFT's

PFT	$V_{cmax,opt}$	$T_{opt}$	$\lambda_{max}$	$Z_{root}$	$\alpha_{leaf}$	$h$	$A_c$	$T_s$	$H_s$
TrBE	50	37	10	1.25	0.12	25	910	-	0.3
TrBR	60	37	10	1.25	0.14	25	180	-	0.3
TeNE	37.5	27	5	1.	0.14	15	910	-	-
TeBE	37.5	32	5	1.25	0.14	15	730	-	-
TeBS	37.5	28	5	1.25	0.14	15	180	12.5	-
BoNE	37.5	25	4.5	1.	0.14	10	910	-	-
BoBS	37.5	25	4.5	1.	0.14	10	180	5	-
BoNS	35	25	4	1.25	0.14	10	180	7	-
NC3	70	$27.5 + 0.25T_l$	2.5	0.25	0.20	0.2	120	4	0.2
NC4	70	36	2.5	0.25	0.20	0.2	120	5	0.2
AC3	90	$27.5 + 0.25T_l$	6	0.25	0.18	0.4	150	10	0.2
AC4	90	36	3	0.25	0.18	0.4	120	10	0.2


# Response to environmental conditions




Temperature



Light



[CO2]



[CO2]

# Concept of externalization

- By default 13 PFT's (named Metaclass) with pre-defined parameters setting
- Most of the parameters can be modified by the user (see <http://forge.ipsl.jussieu.fr/orchidee/wiki/Documentation/OrchideeParameters> or orchidee.default file in the config/PARAM directory)
- The number of PFT's can be extended
  - By setting the NVM parameter and PFT\_TO\_MTC (correspondance array linking a PFT to MTC)




# Outline

---

- A brief history of ORCHIDEE & motivations
- Formalism
- **Main processes**
- Configurations & Inputs requirements



# Main processes



# Resistance terms & Energy budget

## *diffuco module*

- vbeta1 : sublimation
- vbeta2 : interception loss
- vbeta3 : transpiration
- vbeta4 : bare soil evaporation
- vbeta5 : flood plains

## *enerbil module*

- Calculation of :
  - Net radiation
  - Sensible heat flux
  - Latent heat flux
    - Transpiration
    - Evaporation of bare soil
      - and leaf water
    - Sublimation
  - Soil and surface temperature





# Soil temperatures


---

## *thermosoil module*

- Calculates the soil temperatures by solving the heat diffusion equation within the soil
  - the soil is divided into several layers, reaching at least 10m down within the soil. The user can adapt the model to the application.
  - Thickness follows a geometric series.




- Physically-based description of soil water fluxes using Richards equation : 2m soil discretized in at least 11-layers.
- Hydraulic properties based on van Genuchten-Mualem formulation
- Related parameter based on texture (fine, medium, coarse)
- Surface runoff =  $P - E_{sol} - I$
- Free drainage at the bottom



# Vertical discretization in the soils

ORCHIDEE used to have different vertical discretizations for moisture and temperature. The physics require different numerical choices !

This was not tenable any more with soil freezing processes, permafrost and complex snow schemes.



- Users are now provided with a set of parameters to configure the soils.


- zmax\_t (DEPTH\_MAX\_T = 10 )
- zmax\_h (DEPTH\_MAX\_H = 2 )
- depth\_topthickness ( ~1 mm)
- refinebottom
- ratio\_geom\_below




# Routing / Irrigation

*routing module*

- Routing parametrization to calculate water discharge to river



From Guimberteau (thesis, 2010)



The 50 major river basins on the LMD-GCM grid

# Biomass and soil pools

- 9 pools of living biomass
  - Leaves, fine roots, above and below sapwood, above and below heartwood, 'fruits' and short- and long-term 'reserves'
- 4 pools of litter
  - Above/below, Structural & Metabolic
- 3 pools of soil
  - Active, Slow and Passive

x2 Carbon  
Nitrogen

# C assimilation/stomatal conductance

*diffuco module: diffuco\_trans\_co2 routine*

- A and  $G_s$  are calculated at each LAI level:
- Beer-Lambert decrease of light in the canopy
- N-limitation of assimilation:

$$N_L = f(N_{leaf})$$


$$N_L = \frac{k_N \times N_{leaf}}{1 - \exp^{-k_N \times LAI_{Lc}}} \times \exp^{-k_N \times LAI_{Lc}}$$

With  $k_N$  values around 0.1-0.2 (Carrswell et al., 2000; Dewar et al. 2012)

$N_{leaf}$  : leaf nitrogen content  $m^{-2}_{[ground]}$

$N_L$  : leaf nitrogen content  $m^{-2}_{[leaf]}$  at level L

From the leaf to canopy



# Photosynthesis


*diffuco module: diffuco\_trans\_co2 routine*

- Based on Farquahr model
- $V_{c_{max}}$  : photosynthetic capacity ( $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ )

$$V_{c_{max}} = NUE \times N_L$$

with  $NUE$  the Nitrogen Use Efficiency (PFT-dependant) and  $N_L$  the leaf N content ( $\text{gN m}^{-2}_{[\text{leaf}]}$ )

Vmax vs. Leaf N content




Kattge et al. (2009)



# Soil mineral N pools


- Based on the DNDC model (Li et al., 1992, 2000).
- It accounts for:
  - Inputs of mineral through
    - mineralisation
    - N deposition
    - N fertilizers
    - Biological nitrogen fixation
  - Emissions of  $\text{NH}_3$ ,  $\text{NO}$ ,  $\text{N}_2\text{O}$ ,  $\text{N}_2$  by Nitrification and denitrification processes
  - Loss of soil mineral N through
    - Plant N uptake
    - Leaching






# Plant N uptake

- Based on the experimental work of Kronzucker et al. (1995, 1996)



$$N_{\text{up}} = v_{\text{max}} \times N_{\text{min}} \times \left( k_{N\text{min}} + \frac{1}{N_{\text{min}} + K_{N\text{min}}} \right) \times f(T) \times f(\text{NC}_{\text{plant}}) \times C_{\text{root}}$$

Michaelis-Menten type function



Temperature

Fine root mass

N uptake increases in N starved roots

# Allocation of assimilates

*stomate\_growth\_fun\_all module*


- Functional allocation
  - Allometric relationship between sapwood, leaf and root biomass pools
  - Based on Forestry allocation scheme (Dhote and Deleuze)
- N allocation is function of
  - Allocation scheme for Carbon
  - N availability:
    - Leaf C/N ratio is a key variable
    - Varies across two constrained boundaries :  $CN_{\text{leaf,min}}$  and  $CN_{\text{leaf,max}}$



# Allocation of assimilates

*stomate\_growth\_fun\_all module*

- Functional allocation
  - Allometric relationship between sapwood, leaf and root biomass pools
  - Based on Forestry allocation scheme (Dhote and Deleuze)
- N allocation is function of
  - Allocation scheme for Carbon
  - N availability:
    - Leaf C/N ratio is a key variable
    - Varies across two constrained boundaries :  $CN_{\text{leaf,min}}$  and  $CN_{\text{leaf,max}}$



# Phenology

---

*stomate\_phenology module*

- Bud-burst model (Botta et al. 2000)
  - Defined for each PFT based on Growing degree days, Number of chilling days, soil water, ...
  - Calibrated at global scale from bud-burst estimated by satellite

*stomate\_turnover module*

- Senescence
  - Function of leaf age and environmental conditions
  - For trees, a senescence stage is considered until all leaves fall (while for grass senescence it is a continuous process)

# Respirations

- Autotrophic respiration


- Maintenance *stomate\_resp module*

- linear response to temperature with potential adaptation to long-term temperature

- function of Nitrogen content


- Heterotrophic respiration

*stomate\_litter & stomate\_soilcarbon*



- Growth *stomate\_growth\_fun\_all*

- a fixed part of assimilates




# Land-use and land-use change

\* stomate\_lcchange module  
 lcchange + wood\_use routines  
 \* stomate\_woodharvest module

**Wood Harvest**

**Crop Harvest**

45 % of NPP is harvested and respired within the year



SOM decomposition rate of crop is 20% higher

**Only net LUC**

3 woody pools

- 1-yr lifespan
- 10-yr lifespan
- 100-yr lifespan

Fixed fractions over time



# Vegetation dynamic

---

*lpj\_kill, lpj\_pftinout, lpj\_constraints modules*

- Taken from LPJ model
- All PFT's are able to growth in each grid cell
  - Climate constraints define regeneration and adaptation of PFT's
  - Light competition when canopy closure (PFT with NPPmax dominate)
  - Trees always dominate grasses




# Outline

---


- A brief history of ORCHIDEE & motivations
- Formalism
- Main processes
- **Configurations & Inputs requirements**



# Tasks performed by ORCHIDEE



# Infrastructure surrounding ORCHIDEE




- Some numbers to ORCHIDEE's code :
- 83 FORTRAN files
  - 61000 lines of code
  - 43000 lines of comments

# Structure of the code

---

- Use of a modular structure
  - All the variables are dynamics (allocatable)
  - For each module:
    - A main entry point : `<module>_main`
    - An initialisation procedure : `<module>_initialize`
    - An end procedure : `<module>_finalize`
    - An procedure to clear memory : `<module>_clear`
  - All the variables are transmitted by subroutine parameters.
  - Prognostic variables are local to the modules.

# Subroutine Call graph





SECHIBA(30min)  
 STOMATE(30min)  
 STOMATE(1 day)



# Atmospheric Interface

- Meteorological forcing (from monthly to half-hourly)



# Forcing files

---

- Meteorological data
  - One often uses reanalysis or in-situ data with different time resolution (3h, 6h, ½ hour, ...)
  - The spatial resolution of the simulation is driven by the resolution of the meteo forcing file.
  - The time step of a simulation is defined by the parameter TIME\_STEP (30 min by default).
  - The meteorological data often needs to be interpolated in time to the ORCHIDEE time step.

# Ancillary data

---

- Ancillary data needed will depend on the configuration chosen.
- All variables will be interpolated to the grid of ORCHIDEE.
- Some exemples :
  - PFT map and land use
  - Wood harvest intensity
  - Soil texture
  - Soil pH
  - Soil bulk density
  - Background albedo
  - River graphs
  - Topographic slopes
  - Nitrogen deposition
  - Nitrogen fertilisation



# Conclusions

---

- ORCHIDEE is a complex system !
- But you have the chance to use a system which was developed at IPSL and by people who are still present.
- The model has too many options and you will get lost!
- Do not hesitate to ask the original developers if you have problems.
- Enjoy the training !

